arXiv:hep-ph/0502246v3 11 Apr 2006

Probe the effects of split SUSY in rare B decays

Chuan-Hung Chen[1,3*] **and Chao-Qiang Geng**[2,3 †]

[1]*Department of Physics, National Cheng-Kung University, Tainan, 701 Taiwan*

[2]*Department of Physics, National Tsing-Hua University, Hsin-Chu , 300 Taiwan*

[3]*National Center for Theoretical Sciences, Taiwan*

(Dated: February 2, 2008)

Abstract

We study the decays of $B \to K^{(*)}\ell^+\ell^-$ in split supersymmetry with R-parity violation. We find that the decay branching ratio of $B \to K\tau^+\tau^-$ in the new physics model due to the scalar interactions can be 1.8×10^{-6} which is about one order of magnitude larger that in the standard model, whereas those of $B \to K\ell^+\ell^-$ ($\ell = e$ and μ) and the K^* modes are insensitive to the new physics. On the other hand, the forward-backward asymmetries of $B \to K\tau^+\tau^-$ and $K\mu^+\mu^-$, vanishing in the standard model, can be over 10 and 1%, respectively. In addition, we show that the new interactions will significantly change the forward-backward asymmetry in $B \to K^*\tau^+\tau^-$.

---

[*] Email: physchen@mail.ncku.edu.tw

[†] Email: geng@phys.nthu.edu.tw

I. INTRODUCTION

One of the possible extension of the standard model (SM) is supersymmetry (SUSY). It is found that the effects of SUSY at the scale Λ of $O(\text{TeV})$ can solve not only the hierarchy problem, but also the problem of unified gauge coupling [1, 2]. Moreover, the predicted lightest neutralino in supersymmetric models could also provide the candidate of dark matter [1, 3]. In spite of the above successes, models with SUSY still suffer some difficulties from phenomenological reasons, such as the problems on small CP violating phases, large flavor mixings and proton decays, as well as they predict too large cosmological constant. Inevitably, fine tuning always appears in the low energy physics. Recently, in order to explain the cosmological constant problem and preserve the beauty of the ordinary low-energy SUSY models, the scenario of split SUSY is suggested [4, 5], in which the SUSY breaking scale is much higher than the electroweak scale. In this split SUSY scenario, except the SM Higgs which could be as light as the current experimental limit, the scalar particles are all ultra-heavy, denoted by $m_S \sim \mathcal{O}(10^9 - 10^{13})$. On the other hand, by the protection of approximate chiral symmetries, the masses of sfermions, such as gauginos and higgsinos, could be at the electroweak scale [4, 6].

In Ref [7], we have found that due to the large mixings of sneutrinos and the SM-like Higgs, the interesting phenomena on the low energy B_s system, such as the decays of $B_s \to \ell^+\ell^-$ and the $B_s - \bar{B}_s$ mixing, could occur in split SUSY R-parity violating models. It is clear that the same mixings could also give some interesting implications on B_q systems with $q = u$ and d. In this paper, we discuss the possibility to probe the effects of split SUSY in $B \to K^{(*)}\ell^+\ell^-$ decays.

The flavor-changing neutral current (FCNC) processes of $B \to K^{(*)}\ell^+\ell^-$ ($\ell = e, \mu, \tau$) are suppressed and induced by electroweak penguin and box diagrams in the SM with decay branching ratios (BRs) of $\mathcal{O}(10^{-7} - 10^{-6})$ [8, 9, 10]. The decay modes of $B \to K^{(*)}\ell^+\ell^-$ ($\ell = e, \mu$) have been observed by BELLE [11] with $Br(B \to K\ell^+\ell^-) = (5.50^{+0.75}_{-0.70} \pm 0.27 \pm 0.02) \times 10^{-7}$ and $Br(B \to K^*\ell^+\ell^-) = (16.5^{+2.3}_{-2.2} \pm 0.9 \pm 0.4) \times 10^{-7}$, and by BABAR [12] with $Br(B \to K\ell^+\ell^-) = (6.5^{+1.4}_{-1.3} \pm 0.4) \times 10^{-7}$ and $Br(B \to K^*\ell^+\ell^-) = (8.8^{+3.3}_{-2.9} \pm 1.0) \times 10^{-7}$. In addition, BELLE has also reported the forward-backward asymmetry (FBA) as a function of dilepton invariant mass in $B \to K^*\ell^+\ell^-$ [11]. Since the inclusive process of $b \to s\ell^+\ell^-$ arises from loop corrections, these exclusive FCNC rare decays are important for not only testing the SM but also probing new physics. For instance, by considering polarized leptons, various polarization asymmetries have been proposed [13]. On the other hand, in terms of the K^* polarizations ϵ_{K^*}, we can define a triple product operator $\vec{\epsilon}_{K^*} \cdot (\vec{p}_{K^*} \times \vec{p}_\ell)$ to display the CP violating effects under CPT theorem [14, 15]. Moreover, by considering the decaying chain $K^* \to K\pi$, we can investigate various asymmetric operators by studying the angular distributions of the π meson [14]. In the following analysis, we will concentrate on BRs and FBAs.

The paper is organized as follows. In Sec. II, we first present the mechanism to generate the process $b \to s\ell^+\ell^-$ in the scenario of split SUSY. We then derive the differential decay rates and FBAs with the new interactions. In Sec. III, by combining the results of $Br(B_s \to \mu^+\mu^-)$ and the solar neutrino mass, we estimate the decay rates and FBAs of $B \to K^{(*)}\ell^+\ell^-$. Finally, we give our conclusions in Sec. IV.

II. DIFFERENTIAL DECAY RATES FOR $B \to K^{(*)}\ell^+\ell^-$

A. $b \to s\ell^+\ell^-$ in split SUSY without R-parity

In ordinary split supersymmetric models, due to the suppression of the high SUSY breaking scale, one expects that there will be no interesting contributions to low energy physics. Therefore, we extent our consideration to the framework of split SUSY with R-parity violation, in which the conservations of lepton and baryon numbers are broken. For simplicity, in this paper we only consider the lepton number violating effects. The bilinear and trilinear terms for the lepton number violation in the superpotential are written as [16, 17]

$$W = \mu H_1 H_2 + \epsilon_i \mu L_i H_2 + \lambda'_{ijk} L_i Q_j D^c_k + \lambda_{ijk} L_i L_j E^c_k, \tag{1}$$

and the relevant scalar potential is given by

$$V = B H_1 H_2 + B_i L_i H_2 + m^2_{L_i H_1} L_i H^\dagger_1 + h.c. \, . \tag{2}$$

Note that, we have used the same notations for superfields and ordinary fields. The soft parameters B, B_i and $m^2_{L_i H_1}$ could be the same order as the SUSY breaking scale. It has been shown that to solve the atmospheric neutrino mass $\sqrt{\Delta m^2_{atom}} \sim 0.05$ eV, the bilinear related parameters of $\xi_i = m^2_{L_i H_1}/m^2_{L_i} + B_i/m^2_{L_i} \tan\beta - \epsilon_i$ are limited to $10^{-6}/\cos\beta$ at tree level [17], where $\tan\beta = \langle H_2^0\rangle/\langle H_1^0\rangle$. In order to reach the solar neutrino mass scale of $\sqrt{\Delta m^2_{sol}} \sim 9$ meV by the same bilinear couplings, one has to go to one-loop level. However, in split SUSY the results are suppressed by $1/m^2_{L_i}$. It is found that if the trilinear R-parity violating couplings λ'_{i23} and λ'_{i32} are of order one, the problem could be solved by one-loop corrections [17].

Based on the above discussions, if we regard that $\lambda'_{i23,i32}$ as well as the ratios of the bilinear couplings and m_S^2, i.e., $m^2_{L_iH_1}/m_S^2$ and B_i/m_S^2, are order of unity, we find that $b \to s\ell^+\ell^-$ can occur at tree level shown in Fig. 1, which may not be suppressed. Since in the split SUSY approach, except the SM-like Higgs denoted by h^0 is light, all scalars are extremely heavy, we may simplify the calculations by using $-h^0\sin\alpha\,(h^0\cos\alpha)$ instead of the Higgs $H_1^0(H_2^0)$, where the angle α describes the mixing of two neutral Higgses [18]. From Eqs. (1) and (2), the new effective interactions for $b \to s\ell^+\ell^-$ can be described by



FIG. 1: Tree contribution to $b \to s\ell^+\ell^-$ with the cross representing the mixings between sleptons and Higgs.

$$\mathcal{H}_{\text{new}} = \frac{1}{m_h^2}\left(\frac{gm_\ell}{2m_W}\frac{\sin\alpha}{\cos\beta}\right)\frac{m^2_{L_iH_1}}{m^2_{\tilde{\nu}_i}}\left(\lambda'^*_{i23}\bar{s}P_R + \lambda'_{i32}\bar{s}P_L b\right)\bar{\ell}\,\ell\,, \tag{3}$$

where m_h, m_W and m_ℓ stand for the masses of Higgs, W-boson and lepton, respectively, and $m^2_{L_iH_1}$ are from the mixings between sleptons and Higgs. Our purpose of this study is to examine the influence of Eq. (3) on $B \to K^{(*)}\ell^+\ell^-$ decays.

B. Effective Hamiltonian and form factors for $B \to K^{(*)}$ transitions

In order to include the new interactions for $b \to s\ell^+\ell^-$, we write the effective Hamiltonian with interactions of scalar and pseudoscalar to leptons as

$$\mathcal{H}_{\text{eff}} = \frac{G_F\alpha_{em}\lambda_t}{\sqrt{2}\pi}\left[H_{1\mu}L^\mu + H_{2\mu}L^{5\mu} + S_R\bar{\ell}\ell + S_L\bar{\ell}\ell\right] \tag{4}$$

with

$$\begin{aligned}
H_{1\mu} &= C_9^{\rm eff}(\mu)\bar{s}\gamma_\mu P_L b \, - \frac{2m_b}{q^2}C_7(\mu)\bar{s}i\sigma_{\mu\nu}q^\nu P_R b\,, \\
H_{2\mu} &= C_{10}\bar{s}\gamma_\mu P_L b\,, \\
L^\mu &= \bar{\ell}\gamma^\mu\ell\,, \quad L^{5\mu} = \bar{\ell}\gamma^\mu\gamma_5\ell\,, \\
S_R &= C_R\bar{s}P_R b\,, \quad S_L = C_L\bar{s}P_L b\,,
\end{aligned} \tag{5}$$

where α_{em} is fine structure constant, $\lambda_t = V_{tb}V_{ts}^*$, $C_9^{\rm eff}$ and $C_{7,10}$ are the Wilson coefficients (WCs) with their explicit expressions given in Ref. [21] for the SM, $C_{L(R)}$ are from the new interactions of split SUSY, m_b is the current b-quark mass, q is the momentum transfer and $P_{L(R)} = (1 \mp \gamma_5)/2$. Note that $C_9^{\rm eff}$ has included the long-distance effects of $c\bar{c}$ bound states [22].

To obtain the transition elements of $B \to H\,(H = K,\ K^*)$ with various weak vertices, we parametrize them in terms of the relevant form factors as follows:

$$\begin{aligned}
\langle K(p_2)|V_\mu|\bar{B}(p_1)\rangle &= f_+(q^2)\Big\{P_\mu - \frac{P\cdot q}{q^2}q_\mu\Big\} + \frac{P\cdot q}{q^2}f_0(q^2)\,q_\mu, \\
\langle K(p_2)|T_\mu q^\nu|\bar{B}(p_1)\rangle &= \frac{f_T(q^2)}{m_B+m_K}\Big\{P\cdot q\,q_\mu - q^2 P_\mu\Big\}, \\
\langle K^*(p_2,\epsilon)|V_\mu|\bar{B}(p_1)\rangle &= i\frac{V(q^2)}{m_B+m_{K^*}}\varepsilon_{\mu\alpha\beta\rho}\epsilon^{*\alpha}P^\beta q^\rho, \\
\langle K^*(p_2,\epsilon)|A_\mu|\bar{B}(p_1)\rangle &= 2m_{K^*}A_0(q^2)\frac{\epsilon^*\cdot q}{q^2}q_\mu + (m_B+m_{K^*})A_1(q^2)\Big(\epsilon^*_\mu - \frac{\epsilon^*\cdot q}{q^2}q_\mu\Big) \\
&\quad -A_2(q^2)\frac{\epsilon^*\cdot q}{m_B+m_{K^*}}\Big(P_\mu - \frac{P\cdot q}{q^2}q_\mu\Big), \\
\langle K^*(p_2,\epsilon)|T_{\mu\nu}q^\nu|\bar{B}(p_1)\rangle &= -iT_1(q^2)\varepsilon_{\mu\alpha\beta\rho}\epsilon^{*\alpha}P^\beta q^\rho, \\
\langle K^*(p_2,\epsilon)|T^5_{\mu\nu}q^\nu|\bar{B}(p_1)\rangle &= T_2(q^2)\Big(\epsilon^*_\mu P\cdot q - \epsilon^*\cdot q P_\mu\Big) + T_3(q^2)\epsilon^*\cdot q\Big(q_\mu - \frac{q^2}{P\cdot q}P_\mu\Big),
\end{aligned} \tag{6}$$

where $(V_\mu, A_\mu, T_{\mu\nu}, T^5_{\mu\nu}) = \bar{s}(\gamma_\mu, \gamma_\mu\gamma_5, i\sigma_{\mu\nu}, i\sigma_{\mu\nu}\gamma_5)b$, m_{B,K,K^*} are the meson masses of B, K and K^*, $P = p_B + p_{K^{(*)}}$, $q = p_B - p_{K^{(*)}}$ and $P\cdot q = m_B^2 - m_{K^{(*)}}^2$, respectively. By equation of motion, we can have the transition form factors for scalar and pseudoscalar interactions as

$$\begin{aligned}
\langle K|\bar{s}\,b|\bar{B}\rangle &= \frac{P\cdot q}{(m_s-m_b)}\,f_0, \\
\langle K^*|\bar{s}\gamma_5 b|\bar{B}\rangle &= -\frac{2m_{K^*}}{m_b+m_s}\epsilon^*\cdot q\,A_0\,,
\end{aligned} \tag{7}$$

with m_s being the strange-quark mass.

C. Angular Distributions and Forward-Backward Asymmetries

From the definitions of form factors in Eqs. (6) and (7), the transition amplitudes associated with the interactions in Eq. (4) for $B \to K^{(*)}\ell^+\ell^-$ can be written as

$$\mathcal{M}_K = \frac{G_F \alpha_{em} \lambda_t}{\sqrt{2}\pi} \left[m_{97} \bar{\ell} \not{p}_K \ell + m_{10} \bar{\ell} \not{p}_K \gamma_5 \ell + m_5 \bar{\ell} \gamma_5 \ell + m_0 \bar{\ell} \ell \right] \tag{8}$$

with

$$\begin{aligned} m_{97} &= C_9^{\text{eff}} f_+ + \frac{2m_b}{m_B + m_K} C_7 f_T \,, \\ m_{10} &= C_{10} f_+ \,, \\ m_5 &= m_\ell C_{10} \left(f_+ + \frac{P \cdot q}{q^2} (f_0 - f_+) \right) \,, \\ m_0 &= \frac{C_R + C_L}{2} \frac{P \cdot q}{m_s - m_b} f_0 \,, \end{aligned} \tag{9}$$

and

$$\mathcal{M}_{K^*} = \frac{G_F \alpha \lambda_t}{\sqrt{2}\pi} \left\{ \mathcal{M}_{1\mu} \bar{\ell} \gamma^\mu \ell + \mathcal{M}_{2\mu} \bar{\ell} \gamma^\mu \gamma_5 \ell + \tilde{m}_5 \frac{\epsilon^* \cdot q}{q^2} \bar{\ell} \gamma_5 \ell + \tilde{m}_0 \epsilon^* \cdot q \bar{\ell} \ell \right\} \tag{10}$$

where

$$\begin{aligned} \mathcal{M}_{1\mu} &= i \tilde{m}_{97}^1 \varepsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} p_K^\alpha q^\beta - \tilde{m}_{97}^2 \epsilon_\mu^* + \tilde{m}_{97}^3 \epsilon^* \cdot q p_{K\mu}, \\ \mathcal{M}_{2\mu} &= i \tilde{m}_{10}^1 \varepsilon_{\mu\nu\alpha\beta} \epsilon^{*\nu} p_K^\alpha q^\beta - \tilde{m}_{10}^2 \epsilon_\mu^* + \tilde{m}_{10}^3 \epsilon^* \cdot q p_{K\mu}, \end{aligned} \tag{11}$$

with

$$\begin{aligned} \tilde{m}_{97}^1 &= \frac{V}{m_B + m_{K^*}} C_9^{\text{eff}} + \frac{2m_b}{q^2} C_7 T_1 \,, \\ \tilde{m}_{97}^2 &= \frac{1}{2} (m_B + m_{K^*}) C_9^{\text{eff}} A_1 + \frac{1}{2} \frac{2m_b}{q^2} P \cdot q C_7 T_2 \,, \\ \tilde{m}_{97}^3 &= \frac{A_2}{m_B + m_{K^*}} C_9^{\text{eff}} + \frac{2m_b}{q^2} C_7 \left(T_2 + \frac{q^2}{P \cdot q} T_3 \right) \,, \\ \tilde{m}_{10}^i &= \tilde{m}_{97}^i |_{C_7 \to 0, C_9^{eff} \to C_{10}} \,, \\ \tilde{m}_5 &= m_\ell C_{10} \left[-2m_{K^*} A_0 + (m_B + m_{K^*}) A_1 - \frac{q^2 + P \cdot q}{m_B + m_{K^*}} A_2 \right] \,, \\ \tilde{m}_0 &= (C_L - C_R) \frac{m_{K^*}}{m_b + m_s} A_0 \,. \end{aligned} \tag{12}$$

To get the decay rate distributions in terms of the dilepton invariant mass q^2 and the lepton polar angle θ, we use the q^2 rest frame in which $p_\ell = (E_\ell, |\vec{p}_\ell| \sin\theta, 0, |\vec{p}_\ell| \cos\theta)$, $p_H =$

$(E_H, 0, 0, |\vec{p}_H|\cos\theta)$ with $E_\ell = \sqrt{q^2}/2$, $|\vec{p}_\ell| = \sqrt{E_\ell^2 - m_\ell^2}$, $E_H = (m_B^2 - q^2 - m_H^2)/(2\sqrt{q^2})$ and $|\vec{p}_H| = \sqrt{E_H^2 - m_H^2}$. By squaring the transition amplitude in Eq. (8) and including the three-body phase space factor, the differential decay rates as functions of q^2 and θ for $B \to K\ell^+\ell^-$ are given by

$$\begin{aligned}
\frac{d\Gamma_K}{dq^2 d\cos\theta} &= \frac{G_F^2 \alpha_{em}^2 |\lambda_t|^2}{2^8 m_B^2 \pi^5} \sqrt{1 - \frac{(2m_\ell)^2}{q^2}}\, \tilde{p}_K \\
&\times \left[|\vec{p}_K|^2 \left(q^2 - 4|\vec{p}_\ell|^2 \cos^2\theta\right) \left(|m_{97}|^2 + |m_{10}|^2\right) + 4m_K^2 m_\ell^2 |m_{10}|^2 \right. \\
&+ q^2 \left(|m_5|^2 + |m_0|^2 \left(1 - \frac{(2m_\ell)^2}{q^2}\right)\right) + 4m_\ell q \cdot p_K Re(m_{10} m_5^*) \\
&+ \left. 8m_\ell |\vec{p}_K||\vec{p}_\ell| Re(m_{97} m_0^*) \cos\theta \right]. \qquad (13)
\end{aligned}$$

For $B \to K^*\ell^+\ell^-$ decays, by summing up the polarizations of K^* with the identity $\sum \epsilon_\mu^*(p)\epsilon_\nu(p) = (-g_{\mu\nu} + p_\mu p_\nu/p^2)$, from Eq. (10) the differential decay rates are found to be

$$\begin{aligned}
\frac{d\Gamma_{K^*}}{dq^2 d\cos\theta} &= \frac{G_F^2 \alpha_{em}^2 |\lambda_t|^2}{2^8 m_B^2 \pi^5} \sqrt{1 - \frac{(2m_\ell)^2}{q^2}}\, \tilde{p}_{K^*} \left\{ q^2 |\vec{p}_{K^*}|^2 \left(q^2 + 4|\vec{p}_\ell|^2 \cos^2\theta\right) \left(|\tilde{m}_{97}^1|^2 + |\tilde{m}_{10}^1|^2\right) \right. \\
&+ 4m_\ell^2 q^2 |\vec{p}_{K^*}|^2 \left(|\tilde{m}_{97}^1|^2 - |\tilde{m}_{10}^1|^2\right) + \frac{|\vec{p}_{K^*}|^2}{m_{K^*}^2} \left(q^2 - 4|\vec{p}_\ell|^2 \cos^2\theta\right) \left(|\tilde{m}_{97}^2|^2 + |\tilde{m}_{10}^2|^2\right) \\
&+ 2q^2 \left[\left(1 + \frac{2m_\ell^2}{q^2}\right) |\tilde{m}_{97}^2|^2 + \left(1 - \frac{(2m_\ell)^2}{q^2}\right) |\tilde{m}_{10}^2|^2 \right] \\
&+ \frac{q^2}{m_{K^*}^2} |\vec{p}_{K^*}|^4 \left(q^2 - 4|\vec{p}_\ell|^2 \cos^2\theta\right) \left(|\tilde{m}_{97}^3|^2 + |\tilde{m}_{10}^3|^2\right) + 4m_\ell^2 q^2 |\vec{p}_{K^*}|^2 |\tilde{m}_{10}^3|^2 \\
&- 2\frac{q \cdot p_{K^*}}{m_{K^*}^2} |\vec{p}_{K^*}|^2 \left(q^2 - 4|\vec{p}_\ell|^2 \cos^2\theta\right) \left(Re(m_{97}^2 m_{97}^{3*}) + Re(m_{10}^2 m_{10}^{3*})\right) \\
&+ \frac{|\vec{p}_{K^*}|^2}{m_{K^*}^2} \left[|\tilde{m}_5|^2 + q^4 \left(1 - \frac{(2m_\ell)^2}{q^2}\right) |\tilde{m}_0|^2 - 4m_\ell \left(Re\left(\tilde{m}_{10}^2 - \tilde{m}_{10}^3 q \cdot p_{K^*}\right) \tilde{m}_5^*\right) \right] \\
&- 8|\vec{p}_{K^*}||\vec{p}_\ell| q^2 \left[Re(\tilde{m}_{97}^1 m_{10}^{2*}) + Re(\tilde{m}_{97}^2 m_{10}^{1*}) \right] \cos\theta \\
&- \left. 8m_\ell \frac{|\vec{p}_{K^*}||\vec{p}_\ell|}{m_{K^*}^2} \left[q \cdot P_{K^*} Re(\tilde{m}_{97}^2 \tilde{m}_0^*) - q^2 |\vec{p}_{K^*}|^2 Re(\tilde{m}_{97}^3 \tilde{m}_0^*) \right] \cos\theta \right\}. \qquad (14)
\end{aligned}$$

Note that, to have general formulas, we have kept the lepton mass effects in Eqs. (13) and (14). The $\tilde{p}_H$ in both differential decay rates are the spatial momentum of the H meson in the B-meson rest frame, defined as $\tilde{p}_H = \sqrt{E'^2 - m_H^2}$ with $E' = (m_B^2 + m_H^2 - q^2)/(2m_B)$.

By the angular distributions, we can define interesting physical observables, such as the FBAs, given by

$$\mathcal{A}_H^{FB} = \frac{-\int_0^{\pi/2} \frac{d\Gamma_H}{dq^2 d\cos\theta} d\cos\theta + \int_{\pi/2}^{\pi} \frac{d\Gamma_H}{dq^2 d\cos\theta} d\cos\theta}{\int_0^{\pi/2} \frac{d\Gamma_H}{dq^2 d\cos\theta} d\cos\theta + \int_{\pi/2}^{\pi} \frac{d\Gamma_H}{dq^2 d\cos\theta} d\cos\theta}. \qquad (15)$$

It was pointed out that the FBAs of $B \to K^* \ell^+ \ell^-$ are associated with the difference between transverse polarizations of K^* [14], denoted simply by $Re(\mathcal{M}_1^+ \mathcal{M}_2^+) - Re(\mathcal{M}_1^- \mathcal{M}_2^-)$ with

$$\begin{aligned} \mathcal{M}_1^\pm &= \pm |\vec{p}_{K^*}| \sqrt{q^2} \tilde{m}_{97}^1 + \tilde{m}_{97}^2 \,, \\ \mathcal{M}_2^\pm &= \pm |\vec{p}_{K^*}| \sqrt{q^2} \tilde{m}_{10}^1 + \tilde{m}_{10}^2 . \end{aligned} \tag{16}$$

Clearly, the resultant of $Re(\tilde{m}_{97}^1 m_{10}^{2*}) + Re(\tilde{m}_{97}^2 m_{10}^{1*})$ is expected in the SM. For $B \to K \ell^+ \ell^-$, since there are no transverse degrees of freedom, we can infer that there are no FBAs in the SM. Therefore, nonzero FBAs for $B \to K \ell^+ \ell^-$ can be strong evidences of new physics and they have to be from the longitudinal parts. Indeed, from Eq. (13), it is understood that FBA terms are related to $Re(m_{97} m_0^*)$, in which m_0^* is induced from the new scalar interactions. In addition, since the associated FBAs are the interference effects of vector and scalar currents, the mass factor m_ℓ also appears to get the correct chirality.

III. NUMERICAL CALCULATIONS

A. Constraints from the decay of $B_s \to \mu^+ \mu^-$ and the mass of solar neutrino

From Eq. (3), we first consider the $B_s \to \ell^+ \ell^-$ decay, which contains few hadronic uncertainties. The corresponding decay amplitude is given by

$$A = \langle \ell^+ \ell^- | H_{\rm new} | \bar{B}_s \rangle = \frac{-i}{2m_h^2} \left(\frac{g m_\ell}{2 m_W} \frac{\sin\alpha}{\cos\beta} \right) \frac{f_{B_s} m_{B_s}^2}{(m_b + m_s)} \frac{m_{L_i H_1}^2 (\lambda_{i23}^{\prime *} - \lambda_{i32}^\prime)}{m_{\tilde{\nu}_i}^2} \, \bar{\ell} \, \ell , \tag{17}$$

where we have used the identity $\langle 0 | \bar{s} \gamma_5 b | \bar{B}_s \rangle \approx -i \, f_{B_s} m_{B_s}^2 / (m_b + m_s)$ with f_{B_s} being the decay constant. Since the trilinear couplings in sleptons and quarks involve two possible chiralities, there is a cancelation in Eq. (17). Note that if $\lambda_{i23}^{\prime *} = \lambda_{i32}^\prime$, our mechanism vanishes automatically. By including the phase space factor, the decay rate is given by

$$\Gamma \;=\; \frac{m_{B_s}}{16\pi} \frac{G_F m_\ell^2}{\sqrt{2}} \left(\frac{f_{B_s} m_{B_s}}{m_h^2} \frac{\sin\alpha}{\cos\beta} |\mathcal{N}_i| \right)^2 \left[1 - \left(\frac{2 m_\ell}{m_{B_s}} \right)^2 \right]^{\frac{3}{2}} \tag{18}$$

with $\mathcal{N}_i = m_{L_i H_1}^2 (\lambda_{i23}^{\prime *} - \lambda_{i32}^\prime) / m_{\tilde{\nu}_i}^2$. In the SM, it is known that $B_s \to \ell^+ \ell^-$ arises from the electroweak penguin and box diagrams. The decay BR of $B_s \to \mu^+ \mu^-$ is found to be $(3.8 \pm 1.0) \times 10^{-9}$ [19] which is much less than the current experimental upper limit of 5.0×10^{-7} [20]. In order to further limit the values of unknown parameters, we include the solar neutrino mass which is presented by [17]

$$M^{\nu}_{ij} \sim \frac{3}{8\pi^2}\lambda'_{i23}\lambda'_{j32}\frac{m_b m_s}{m_S}. \tag{19}$$

To preserve the solar neutrino mass to be $\sim$ 9 meV, we choose $\lambda'_{i23} = 0.9$, $\lambda'_{i32} = 0.5$, $m_b = 4.5$ GeV, $m_s = 0.13$ GeV, and $m_S = 10^9$ GeV. Since the remaining free parameters are $m_{L_iH_1}/m_{\tilde{\nu}_i}$ and m_h, to illustrate our numerical results, we set $m_{L_iH_1}/m_{\tilde{\nu}_i} = 0.2$ and $m_h = 150$ GeV and take $m_{B_s} = 5.37$ GeV, $f_{B_s} = 0.23$ GeV, $\tau_{B_s} = 1.46 \times 10^{-12}\, s$ and $\alpha = \pi/2 + \beta$. As a result, we get $Br(B_s \to \mu^+\mu^-) = 1.66 \times 10^{-8}$ which is one order of magnitude larger than that predicted in the SM. It is clear that a heavier Higgs or a smaller $m_{L_iH_1}/m_{\tilde{\nu}_i}$ will make the contribution be smaller. In the following numerical calculations, we will adopt the above values of parameters to estimate BRs and FBAs in $B \to K^{(*)}\ell^+\ell^-$ decays.

B. Decay rate distributions and branching ratios

By integrating the lepton polar angle, the differential decay rate as a function of the dilepton invariant mass for $B \to K\ell^+\ell^-$ is given by

$$\begin{aligned}
\frac{d\Gamma_K}{dq^2} &= \frac{G_F^2\alpha_{em}^2|\lambda_t|^2}{2^8 m_B^2 \pi^5}\sqrt{1-\frac{(2m_\ell)^2}{q^2}}\,\tilde{p}_K \\
&\times \left[\frac{4}{3}|\vec{p}_K|^2\left(q^2+2m_\ell^2\right)\left(|m_{97}|^2+|m_{10}|^2\right) + 8m_K^2 m_\ell^2|m_{10}|^2\right. \\
&+ \left. 2q^2\left(|m_5|^2+|m_0|^2\left(1-\frac{(2m_\ell)^2}{q^2}\right)\right) + 8m_\ell q\cdot p_K Re(m_{10}m_5^*)\right].
\end{aligned} \tag{20}$$

Similarly, for $B \to K^*\ell^+\ell^-$, one has

$$\begin{aligned}
\frac{d\Gamma_{K^*}}{dq^2} &= \frac{G_F^2\alpha_{em}^2|\lambda_t|^2}{2^8 m_B^2 \pi^5}\sqrt{1-\frac{(2m_\ell)^2}{q^2}}\,\tilde{p}_{K^*}\left\{\frac{8}{3}q^2|\vec{p}_{K^*}|^2\left(q^2-m_\ell^2\right)\left(|\tilde{m}^1_{97}|^2+|\tilde{m}^1_{10}|^2\right)\right. \\
&+ 8m_\ell^2 q^2|\vec{p}_{K^*}|^2\left(|\tilde{m}^1_{97}|^2-|\tilde{m}^1_{10}|^2\right) + \frac{4}{3}\frac{|\vec{p}_{K^*}|^2}{m_{K^*}^2}\left(q^2+2m_\ell^2\right)\left(|\tilde{m}^2_{97}|^2+|\tilde{m}^2_{10}|^2\right) \\
&+ 4q^2\left[\left(1+\frac{2m_\ell^2}{q^2}\right)|\tilde{m}^2_{97}|^2+\left(1-\frac{(2m_\ell)^2}{q^2}\right)|\tilde{m}^2_{10}|^2\right] \\
&+ \frac{4}{3}\frac{q^2}{m_{K^*}^2}|\vec{p}_{K^*}|^4\left(q^2+2m_\ell^2\right)\left(|\tilde{m}^3_{97}|^2+|\tilde{m}^3_{10}|^2\right) + 8m_\ell^2 q^2|\vec{p}_{K^*}|^2|\tilde{m}^3_{10}|^2 \\
&- \frac{8}{3}\frac{q\cdot p_{K^*}}{m_{K^*}^2}|\vec{p}_{K^*}|^2\left(q^2+2m_\ell^2\right)\big(Re(m^2_{97}m^{3*}_{97}+Re(m^2_{10}m^{3*}_{10})\big) + 2\frac{|\vec{p}_{K^*}|^2}{m_{K^*}^2} \\
&\times \left.\left[|\tilde{m}_5|^2+q^4\left(1-\frac{(2m_\ell)^2}{q^2}\right)|\tilde{m}_0|^2-4m_\ell\left(Re\left(\tilde{m}^2_{10}-q\cdot p_{K^*}\tilde{m}^3_{10}\right)\tilde{m}_5^*\right)\right]\right\}.
\end{aligned} \tag{21}$$

To calculate the numerical values, we use $m_B = 5.28$ GeV, $m_K = 0.5$ GeV, $m_{K^*} = 0.89$ GeV, $\alpha = 1/129$, $V_{ts} = 0.40 \pm 0.003$, $\tau_{B^0} = 1.536 \times 10^{-12}$ s and $\tau_{B^+} = 1.671 \times 10^{-12}$ s. For the form factors in $\langle K^{(*)}|\Gamma|\bar{B}\rangle$, we quote the updated results of the light-cone sum rules (LCSR) [23]. The explicit fitting forms are summarized as

$$\begin{aligned}
f_+(q^2) &= \frac{0.1903}{1-q^2/29.3} + \frac{0.1478}{(1-q^2/29.3)^2}, \quad f_0(q^2) = \frac{0.3338}{1-q^2/38.98}, \\
f_T(q^2) &= \frac{0.1851}{1-q^2/29.3} + \frac{0.1905}{(1-q^2/29.3)^2}, \\
V(q^2) &= \frac{0.923}{1-q^2/5.32^2} - \frac{0.511}{1-q^2/49.4}, \quad A_0(q^2) = \frac{1.364}{1-q^2/5.282^2} - \frac{0.99}{1-q^2/36.78}, \\
A_1(q^2) &= \frac{0.29}{1-q^2/40.38}, \quad A_2(q^2) = -\frac{0.084}{1-q^2/52.} + \frac{0.342}{(1-q^2/52.)^2}, \\
T_1(q^2) &= \frac{0.823}{1-q^2/5.32^2} - \frac{0.491}{1-q^2/46.31}, \quad T_2(q^2) = \frac{0.333}{1-q^2/41.41}, \\
T_3(q^2) &= -\frac{0.036}{1-q^2/48.1} + \frac{0.368}{(1-q^2/48.1)^2}\,. \qquad (22)
\end{aligned}$$

In order to exclude the backgrounds from $B \to J/\Psi(\Psi')K^{(*)}$, we follow the BELLE's veto windows [11], defined as:

$$\begin{aligned}
\mathrm{I_V}: &\quad -0.20\mathrm{GeV} < M_{e^+e^-} - m_V < 0.07\mathrm{GeV}, \\
\mathrm{II_V}: &\quad -0.10\mathrm{GeV} < M_{\mu^+\mu^-} - m_V < 0.08\mathrm{GeV} \qquad (23)
\end{aligned}$$

for $B \to K\ell^+\ell^-(\ell = e,\, \mu)$ with $V = \Psi,\, \Psi'$ and for $B \to K^*\ell^+\ell^-$ $(\ell = e, \mu)$ with $V = \Psi'$, and

$$\begin{aligned}
\mathrm{III_\Psi}: &\quad -0.25\mathrm{GeV} < M_{e^+e^-} - m_{J/\Psi} < 0.07\mathrm{GeV}, \\
\mathrm{IV_\Psi}: &\quad -0.15\mathrm{GeV} < M_{\mu^+\mu^-} - m_{J/\Psi} < 0.08\mathrm{GeV} \qquad (24)
\end{aligned}$$

for $B \to K^*\ell^+\ell^-(\ell = e,\, \mu)$ decays. For $B \to K^{(*)}\tau^+\tau^-$, we use

$$V_{\Psi'}: \quad M_{\tau^+\tau^-} - m_{\Psi'} < 0.08\,. \qquad (25)$$

Hence, the predicted values in the SM for various lepton modes are presented in Table I. Note that, the values in the table are obtained by taking the average lifetime in charged and neutral B mesons. Furthermore, according to the results of Table I, the average BRs for the $\ell = e$ and μ modes in the SM are given by

$$\begin{aligned}
Br(B \to K\ell^+\ell^-) &= 5.89 \times 10^{-7}\,, \\
Br(B \to K^*\ell^+\ell^-) &= 13.67 \times 10^{-7}\,. \qquad (26)
\end{aligned}$$

TABLE I: The average BRs (in units of 10^{-7}) of $B \to K^{(*)}\ell^+\ell^-$ for charged and neutral B decays in the SM with the veto windows defined in Eqs. (23), (24) and (25).

Mode	$B \to Ke^+e^-$	$B \to K\mu^+\mu^-$	$B \to K\tau^+\tau^-$
BR	5.59	6.18	1.46
Mode	$B \to K^*e^+e^-$	$B \to K^*\mu^+\mu^-$	$B \to K^*\tau^+\tau^-$
BR	14.79	12.54	1.75

In the split SUSY R-parity violating model, we obtain

$$\begin{aligned} Br(B \to K\ell^+\ell^-) &= 6.14 \times 10^{-7}\,, \\ Br(B \to K\tau^+\tau^-) &= 17.95 \times 10^{-7}\,, \\ Br(B \to K^*\ell^+\ell^-) &= 13.87 \times 10^{-7}\,, \\ Br(B \to K^*\tau^+\tau^-) &= 1.91 \times 10^{-7}. \end{aligned} \tag{27}$$

It is interesting to see that the decay BR of $B \to K\tau^+\tau^-$ gets one order of magnitude enhancement, while that of $B \to K^*\tau^+\tau^-$ changes only a little. Hence, the new scalar interactions could largely enhance the BR of $B \to K\tau^+\tau^-$, but not that of the K^* mode. In Figs. 2 and 3, we show the differential decay BRs of $B \to K^0\mu^+\mu^-$ and $K^0\tau^+\tau^-$, where the dashed and solid lines stand for the predictions with and without new physics, respectively. Since the scalar interactions are associated with the lepton mass, as displayed by the figures, the influences on light lepton modes are small.



FIG. 2: Differential BRs (in units of 10^{-7}) for (a) $B \to K^0 \mu^+\mu^-$ and (b) $B \to K^0 \tau^+\tau^-$, where the dashed and solid lines correspond to the results with and without new physics, respectively.



FIG. 3: Same as Fig. 2 but for (a) $B \to K^{*0}\mu^+\mu^-$ and (b) $B \to K^{*0}\tau^+\tau^-$.

C. Forward-backward asymmetries

It is known that FBAs of $b \to s\ell^+\ell^-$ decays could be sensitive to new physics [8, 14, 22, 24]. From Eqs. (13), (14) and (15), the FBAs for $B \to K\ell^+\ell^-$ and $K^*\ell^+\ell^-$ are given by

$$\mathcal{A}_K(q^2) = \frac{1}{d\Gamma_K/dq^2}\frac{G_F^2\alpha_{em}^2|\lambda_t|^2}{2^8 m_B^2\pi^5}\sqrt{1-\frac{(2m_\ell)^2}{q^2}}\tilde{p}_K\Big[-8m_\ell|\vec{p}_K||\vec{p}_\ell|Re(m_{97}m_0^*)\Big]\,, \quad (28)$$

$$\begin{aligned}\mathcal{A}_{K^*}(q^2) &= \frac{1}{d\Gamma_{K^*}/dq^2}\frac{G_F^2\alpha_{em}^2|\lambda_t|^2}{2^8 m_B^2\pi^5}\sqrt{1-\frac{(2m_\ell)^2}{q^2}}\tilde{p}_{K^*} \\ &\times\ \Big\{8|\vec{p}_{K^*}||\vec{p}_\ell|q^2\left[Re(\tilde{m}_{97}^1 m_{10}^{2*})+Re(\tilde{m}_{97}^2 m_{10}^{1*})\right] \\ &+\ 8m_\ell\frac{|\vec{p}_{K^*}||\vec{p}_\ell|}{m_{K^*}^2}\left[q\cdot P_{K^*}Re(\tilde{m}_{97}^2\tilde{m}_0^*)-q^2|\vec{p}_{K^*}|^2 Re(\tilde{m}_{97}^3\tilde{m}_0^*)\right]\Big\}\,, \quad (29)\end{aligned}$$

respectively. As discussed before, the new contributions from scalar interactions are suppressed due to the light lepton masses. However, since $\mathcal{A}_K(q^2)$ vanishes in the SM, the search of the FBAs in $B \to K\ell^+\ell^-$ with lots of accumulated B samples is still important in future B factories, such as SuperB, BTeV and LHCB. For this reason, we would display the FBAs of $B \to K\ell^+\ell^-$ ($\ell = \mu$ and τ) associated with the new interactions as functions of q^2 in Fig. 4. Here, we have also used the criterions of Eqs. (23), (24) and (25) to exclude the backgrounds. From the figure, we see clearly that $\mathcal{A}_K(B \to K\ell^+\ell^-)$ can be as large as 1 and 10% for $\ell = \mu$ and τ, which require at least 2×10^{10} and 6×10^{7} B samples for experimental observations at 1σ level. In Fig. 5, we present the FBAs of $B \to K^*\ell^+\ell^-$. From Fig. 5(a), we



FIG. 4: Forward-backward asymmetries (FBAs) for (a) $B \to K\mu^+\mu^-$ and (b) $B \to K\tau^+\tau^-$.



FIG. 5: Forward-backward asymmetries (FBAs) for (a) $B \to K^*\mu^+\mu^-$ and (b) $B \to K^*\tau^+\tau^-$, where dashed and solid lines represent the results with and without new physics, respectively.

find that the new interactions have no effect on $\mathcal{A}_{K^*}(B \to K^*\mu^+\mu^-)$, but a large influence on $\mathcal{A}_{K^*}(B \to K^* \tau^+ \tau^-)$.

IV. CONCLUSIONS

We have studied the decays of $B_s \to \mu^+\mu^-$ and $B \to K^{(*)}\ell^+\ell^-$ in split SUSY with R-parity violation. With the new updated form factors calculated by the light-cone sum rules, we have obtained the decay branching ratios within the BELLE's veto windows to be $Br(B \to K^{(*)}\ell^+\ell^-) = 5.89\ (13.67) \times 10^{-7}$ for $\ell = e$ and μ in the SM. If we set the parameters of new interactions such that the BR of $B_s \to \mu^+\mu^-$ is one order of magnitude larger than that of SM, we find that the decay branching ratio of $B \to K\tau^+\tau^-$ can be 1.8×10^{-6} which is about one order of magnitude larger that in the SM, whereas those of $B \to K\ell^+\ell^-$ ($\ell = e$ and μ) and the K^* modes are insensitive to the new effects. For the FBAs, we have shown that $\mathcal{A}_K$ for $B \to K\tau^+\tau^-$ and $K\mu^+\mu^-$, which are zero in the SM, can be over 10 and 1% in our new physics model, respectively. In addition, we have also demonstrated that the new interactions can significantly change the spectrum of $\mathcal{A}_{K^*}$ in $B \to K^*\tau^+\tau^-$.

Acknowledgments

This work is supported in part by the National Science Council of R.O.C. under Grant #s: NSC-93-2112-M-006-010 and NSC-93-2112-M-007-014.

---

[1] S. Dimopoulos and H. Georgi, Nucl. Phys. B**193**, 150 (1981).

[2] S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D**24**, 1681 (1981).

[3] H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983).

[4] N. Arkani-Hamed and S. Dimopoulos, hep-th/0405159.

[5] N. Arkani-Hamed, S. Dimopoulos, G.F. Giudice and A. Romanino, Nucl. Phys. B**709**,3 (2005).

[6] G.F. Giudice and A. Romanino, Nucl. Phys. B**699**, 65 (2004).

[7] C.H. Chen and C.Q. Geng, hep-ph/0501001.

[8] A. Ali *et al.*, Phys. Rev. D**61**, 074024 (2000).

[9] D. Melikhov, N. Nikitin, and S. Simula, Phys. Rev. D**57**, 6814 (1998).

[10] C.H. Chen and C.Q. Geng, Phys. Rev. D**63**, 114025 (2001).

[11] Belle Collaboration, K. Abe *et al.*, hep-ex/0410006.

[12] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **91**, 221802 (2003).

[13] F. Kruger and L. M. Sehgal, Phys. Lett. B**380**, 199 (1996); J.L. Hewett, Phys. Rev. D**53**, 4964 (1996); C.Q. Geng and C.P. Kao, Phys. Rev. D**54**, 5636 (1996); C.Q. Geng and C.P. Kao, Phys. Rev. D**57**, 4479 (1998); T.M. Aliev *et al.*, Phys. Rev. D**64**, 055007 (2001); W. Bensalam *et al.*, Phys. Rev. D**67**, 034007 (2003); S.R. Choudhury *et al.*, Phys. Rev. D**68**, 054016 (2003);

[14] C.H. Chen and C.Q. Geng, Nucl. Phys. B**636**, 338 (2002); Phys. Rev. D **66**, 014007 (2002).

[15] T.M. Aliev *et al.*, Phys. Rev. D **66**, 115006 (2002).

[16] K. Cheung and O.C.W. Kong, Phys. Rev. D**64**, 095007 (2001).

[17] E.J. Chun and J.D. Park, JHEP **0501**, 009 (2005).

[18] J.F. Gunion *et al.*, *The Higgs Hunter's Guide* (Addison-Wesley, Reading, M.A., 1990).

[19] A.J. Buras, Phys. Lett. B**566**, 115 (2003).

[20] D0 Collaboration, V.M. Abazov *et al.*, Phys. Rev. Lett. **94**, 071802 (2005).

[21] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys **68**, 1230 (1996).

[22] C.H. Chen and C.Q. Geng, Phys. Rev. D**66**, 034006 (2002).

[23] P. Ball and R. Zwicky, Phys. Rev. D**71**, 014015 (2005); Phys. Rev. D**71**, 014029 (2005).

[24] Q.S. Yan *et al.*, Phys. Rev. D**62**, 094023 (2000) ; D.A. Demir *et al.*, Phys. Rev. D**66**, 034015 (2002); A. Arda and M. Boz, Phys. Rev. D**66**, 075012 (2002); T. Feldmann and J. Ma, JHEP **0301**, 074 (2003); S. R. Choudhury *et al.*, Phys. Rev. D**69**, 054018 (2004).